

## NHEXAS Arizona Arizona Border Survey Overview



Michael D. Lebowitz P.I.

Co-Principal Investigators

Mary Kay O'Rourke, Arizona

Sydney M. Gordon, Battelle

Demetrios Moschandreas, IIT




## Overarching Vision and Goal

Vision: To select a representative population and obtain all the measurements needed to perform complete exposure assessment, and to enhance the quality of policy formulation with appropriate use of the survey results.

Primary Goal: To define the high end (upper 10<sup>th</sup> percentile) of the exposure distribution for the population.



## NHEXAS AZ & BORDER AZ





#### Objectives & Hypotheses: NHEXAS AZ

- (1) To document the occurrence, distributions and determinants of exposure.
- (2) To evaluate geographic and temporal trends.
  - exposures detrimental to public health do not occur
  - temporal trends do not vary
  - there are no differences among different geographic areas
- (3) To evaluate the impact on exposure models of using less precise concentration measurements.



#### Objectives & Hypotheses: NHEXAS AZ

- (4) To evaluate exposure as indicated using biomarkers.
- (5) To link "exposure-dose-health" information to enhance, surveillance, risk assessment, risk management and public health policies.
- (6) To compare iterative exposure assessment models (various levels of complexity) and assess the impact of the EA models on risk assessments.
- (7) Evaluate model uncertainties.



#### Objectives & Hypotheses: NHEXAS AZ


- (8) to collaborate on pharmacokenetic modeling.
- (9) To evaluate exposures of sub-populations by modeling.

Issues of Stratification and Detection



#### Objectives & Hypotheses: Border

- The distribution of exposure for the population residing in the "Border" region does/does not differ from that of the State.
- The intermedia analyte relationships do/do not differ between the Border and the rest of the State.
  - Compare AZ Border media means with LRGV border results



### Target Analytes

NHEXAS-- PM<sub>10</sub>

Metals: Pb, As, Ni, Cd, Cr, Mn, Ba, V, Se, Zn (+17)

Pesticides: Chlorpyrifos, Diazinon, Malathion

VOCs: Benzene, Toluene, 1,3 butadiene,

TCE, Formaldehyde (+39)

**Border** -- NHEXAS Analytes plus PM<sub>2.5</sub>

Additional Pesticides--OCs

**PAHs** 



#### Study Design: NHEXAS & Border

- Recruitment: Population Proportional to Size probability of selection is proportional to the number of occupied housing units in the 1990 census.
- Multiple Media & Multiple Pathways
  - Air, Water, Soil, Dust, Wipes
  - Food (Solid & Liquid)
  - Biologicals (Blood, Urine)
  - Questionnaires, including Time-Activity Diaries



#### Recruitment Strata

#### NHEXAS AZ


- Areas = 14/15 Counties
- PSUs = 49 Census Tracts
- SSUs= 245 Block Combos (5 per tract)
- Houses = 5 houses per block


3-4 Stage 3's per PSU


#### AZ Border Survey


- Areas = 3/4 Counties
- PSUs = 25Census Tracts
- SSUs= 100 Block Combos (4 per tract)
- Houses = 3 houses per block


4 Stage 3's per PSU


















#### Study Design

- Three Phase Design:
  - Stage I NHEXAS = 1225 Border = 300

Rate 77.9% ~87.6%

Descriptive Baseline Questionnaires.

- Stage II NHEXAS = 391 Border = ~91 Questionnaires, Food & Activity Diaries & Collection of Screening Data & Samples.
- Stage III NHEXAS = 179 Border = 86
   Intensive Environmental Sampling, Biomarkers, Water
   Duplicate Diet & Questionnaires.



#### Stage 1: Recruitment

- During early recruitment, homes were contacted up to 15 times with no success
- Negligible recruitment occurred after 5 attempts.
- Procedure of NHEXAS AZ & Border
   Weekday, Weekend day, Weekday evening,
   Weekend evening, one other attempt
- Virtually NO ONE would complete a 27 page (1 hr) Baseline QX unscheduled.



#### NHEXAS Demographics: Race/Ethnicity

	White	Black	America Indian	Asian	Other	Hispanic	No Response
% 1990 Census	81.0*	3.0	5.6	1.4	$9.0^*$	18.6	
% 1996 Census Estimate	88.9	2.97	4.99	1.75	NA	21.26	
% Total NHEXAS	92.5	2.1	3.5	0.5	1.2	41.8	0.2
% Primary Stage I	93.3	1.9	2.8	0.5	1.4	35.2	0.1
% Primary Stage II	93.7	1.6	2.2	0.6	2.0	30.1	0.0
% Primary Stage III	91.7	1.7	2.8	0.6	3.3	30.0	0.0



## Demographic Characteristics of Stage III NHEXAS Participants

	<u>Percent</u>	<u>Number</u>	
Age Group			
6-17	19.5%	35	
18-65	65.4%	117	
>65	15.1%	27	
Hispanic Ethnicity			
Yes	29.6%	53	
No	70.4%	126	
Smoking			
Yes	18.9%	34	
No	81.1%	145	



#### NHEXAS & Border AZ Essentials

- Multiethnic, multigender Field Teams
- Bilingual Field Teams
- All materials in Spanish and English
  - NHEXAS:
    - 25 % preferred Spanish
    - 4% completed Questionnaires in Spanish
  - Border:
    - ~ 75% preferred Spanish
    - ~50% completed Questionnaires in Spanish



## Stages 2 & 3: Sampling

- Randomize homes and recruit for Stage 2 & 3 sampling.
- If selected for Stage 3 sampling then collect Stage 2 samples at the same time.
- If sampling is refused, try to obtain a Baseline Questionnaire



#### **Stage 2 (n=125)**

#### Questionnaires:


Descriptive Update
Baseline
Diet Diary\*
Time / Activity\*
Technician
Supplement

\* One day recall

Day 1Day 2Day 3Day 4Day 5Day 6Day 7


#### Sample Collection:

Yard Soil Floor Dust





# NHEXAS AZ Stage III Households





## Households Completed

	<b>NHEXAS</b>	<b>AZ Border</b>
# HH (IRN 01)	955/1225	263/300
	(78%)	(88%)
# People (D Qx)	3205	~833
# Baselines (IRN 01)	525	169
# Secondary Baseline Qx	564	~157
#HH Stage 2 only	212	5
# HH Stage 2 + 3	179	86



#### NHEXAS Field and Lab Methods

	Field	Lab
Air		
PM <sub>10</sub> In & Out	Pump & Teflo Filter 3 of 7 days.	HGAAS (As)
PM <sub>10</sub> Personal Air	Pump & Teflo Filter 8 of 24 hrs.	ICP, XRF
House Dust		
Vacuum Floor	In-line filter, sieve to 62.5 µm.	HGAAS (As)
Window Sill Wipe	Gauze & DiDw wipe.	ICP, XRF
Soil		
Yard	Composite, sieve to 62.5 µm.	HGAAS (As)
Foundation	Composite, sieve to 62.5 µm.	ICP, XRF
Food		
Solid & Beverage	24 hr. Duplicate sample.	ICP-MS by FDA
Water		
Tap	3 min flush (EPA 200.8).	ICP-MS
Drinking	Standing sample(EPA 200.8).	ICP-MS
Dermal Wipe		
Both Hands	Gauze wipe, DiDw x 2 mins.	GCMS
Blood		
(Pb & Cd only)	10 mL Venipuncture	GFAAS by CDC
Urine		
	First Morning Void	HGFAA (As)
		GFAAS



#### NHEXAS Detection Limits

• see overheads



## NHEXAS Distribution of As Concentration by Media

		Detection	Number of samples	% of samples	Range of values above	Percentile		e
Media	Metal	method	evaluated	BDL	MDL	50 <sup>th</sup>	75 <sup>th</sup>	90 <sup>th</sup>
Air—In	As	HG-AAS	125	71	2.9-22.3	BDL	3.5	7.4
Air—Out	As	HG-AAS	116	68	3.1-24.5	BDL	5.3	8.9
Dust	As	HG-AAS	131	0	0.3-50.6	6.7	10.4	16.5
Soil	As	HG-AAS	143	0	1.8-69.3	8.3	12.5	19.2
Food	As	ICP-MS	159	0.6	2.3-2878.0	9.0	15.1	30.6
Beverage	As	ICP-MS	154	30	1.0-19.9	1.9	3.8	6.8
Drinking								
Water								
Consumed	As	ICP-MS	73	59	0.2-15.9	BDL	2.9	6.5
Tap								
Water								
Consumed	As	ICP-MS	82	0	0.6-36.7	4.7	9.1	15.1



### Potential As Dose µg/day

Mining towns (n=43 subjects)

	25th	50th	75th	90th
◆ Dust	2.1	2.8	4.0	5.7
◆ Soil	.5	.8	1.2	1.6
◆ Water	.7	3.3	12.2	15.2
◆ Food	4.5	6.1	9.4	22.6
◆ Beverage	1.5	2.5	7.0	16.6
◆ Total	14.6	20.6	35.8	69.2



#### Potential As Dose µg/day

Non Mining towns (n=122 subjects)

	25th	50th	75th	90th
◆ Dust	1.5	2.0	2.3	3.4
◆ Soil	.3	.5	0.8	1.4
◆ Water	.5	2.6	4.2	8.1
◆ Food	3.2	5.5	10.3	22.0
◆ Beverage	.7	1.8	3.8	6.7
◆ Total	11.6	15.6	23.5	33.6



#### Conclusions

- The last example illustrates the power of these data to identify total exposure during the sample week.
- The As data demonstrates a geographic difference in exposure.
- We are in the final stages of compiling the Border data bases. No work has been done on these data.



#### Conclusions

- The percentage of samples with BDL values are very high for many analytes.
- Still, NHEXAS and affiliated surveys provide rich sources of data for examination.

#### **Data Collection Schedule**

